

PILARIX

Cream

PILARIX CREAM IS A MODERN PRODUCT WHICH EFFECTIVELY CONTROLS THE EXCESSIVE KERATINISATION OF THE SKIN. BY INCREASING THE HYDRATION OF THE STRATUM CORNEUM IT IMPROVES SKIN REGENERATION PROCESSES. PILARIX CREAM HAS SOOTHING EFFECT AND INTENSIVELY MOISTURISES.

Information about the product. Read all of this leaflet carefully before the use.

PILARIX CREAM CONTAINS:

SALICYLIC ACID 2% facilitates the separation of the accumulated thickened stratum corneum
UREA 20% highly moisturising and increases flexibility of stratum corneum

EXCIPIENTS stabilise proper hydration and lubrication of the skin

PANTHENOL has soothing, anti-inflammatory effects in cases of skin irritation, regardless the cause

ALLANTOIN accelerates the regeneration of irritated skin, biologically damaged

SHEA BUTTER provides the skin softness, smoothness, protects from cold, wind and sunlight, enables faster healing of the skin.

PILARIX CREAM IS USED FOR:

Excessively hard, dry skin on elbows, knees, heels
Chronically dry skin
As an ideal mean of toning - soothing after epilation
As a mean of improving skin regeneration

PILARIX CREAM INDICATIONS FOR USE ON SKIN CARE:

Follicular hyperkeratosis
Eczema in excessive flaking phase
Chronically dry skin
As a means of accelerating the regeneration of the skin between the steroid therapy periods

HOW TO USE PILARIX EMULSION:

Apply on a dry skin 2 a day or according to recommendations of a doctor. In a case of skin irritation, one should contact a doctor.

SIDE EFFECTS:

In case of emulsion application on a damaged, irritated, inflamed skin may occur worsening of inflammation accompanied with pruritus. In such case one should stop using the cream.

CONTRAINDICATIONS:

Hypersensitivity to salicylic acid, urea or other ingredients of the cream, pregnancy, lactation period, age under 12 years of age, hepatic, renal disorders. Do not use in the case of bacterial, fungal, viral inflammation. Do not apply the cream on a damaged skin mucous membranes, face, eyes and genitourinary area.

ADDITIONAL RECOMMENDATIONS

AND PRECAUTIONS:

For external use only. If accidentally swallowed, immediately contact a doctor. Keep out of the reach of children. After using the cream, package should be closed. Store in a dry and dark place at room temperature (15°C to 25°C).

The product dermatologically tested. Received a positive assessment of the operational effectiveness on the basis of the research conducted at the Institute of Medical Cosmetics, Swietokrzyska street 33a in Warsaw.

AVAILABLE PACKAGING:

PILARIX cream - tube 50 ml or 100 ml
After opening the usefulness of 6 months.



LEFROSCH

LEFROSCH Sp. z o.o.
ul. Bernardyńska 2, 64-000 Kościan
E-MAIL: BIURO@LEFROSCH.COM

www.lefrosch.com